



DRUGS USED FOR THE TREATMENT OF COVID-19 INFECTION

Last update: [16.03.2020]

LOPINAVIR / RITONAVIR	INTERFERON Beta-1	REMEDSIVIR	HYDROXY-CHLOROQUINE	TOCILIZUMAB
		 <i>Dilution:</i> 100-250 ml NaCl <i>Stability:</i> • 4 h ambient temp. • 24 h refrigerator		 <i>Dilution:</i> 100 ml NaCl <i>Stability:</i> • 4 h ambient temp. • 24 h refrigerator
<ul style="list-style-type: none"> • 200 mg / 50 mg every 12 hours (oral). • Suspension inj.: 5 ml / 12 h. 	<ul style="list-style-type: none"> • 250 mcg (1 vial) / 48 h (SC) 	<ul style="list-style-type: none"> • Dose: 200 mg (IV) day 1 • Maintenance: 100 mg/ 24 h (days 2-10) 	<ul style="list-style-type: none"> • Dose: 400 mg (IV) day 1 • Maintenance: 200 mg/ 12 h (days 2-14) 	<ul style="list-style-type: none"> • 8 mg/kg IV (max. 800mg) • New rate admin. at 12 h.
<p>MAX DURATION: 14 days</p> <p><u>PRECAUTIONS:</u></p> <ul style="list-style-type: none"> • Do not shred/crush tab. • Inhibitors of CYP3A4. • Gastrointestinal A.E • Avoid in insufficiency Severe Liver Disease. ↓ [Fosamprenavir / Ritonavir] 	<p>MAX DURATION: 14 days</p> <p><u>PRECAUTIONS:</u></p> <ul style="list-style-type: none"> • Flu-like syndrome • Avoid in psychological illness / serious depression. 	<p>MAX DURATION: 10 days</p> <p><u>PRECAUTIONS:</u></p> <ul style="list-style-type: none"> • Clean via with 30 mL NaCl after administration. • Admin. In 30-60 minutes. • Monitor possible hypotension infusion. <p> <i>Suspend lopinavir / ritonavir and Interferon B</i></p>	<p>MAX DURATION: 14 days</p> <p><u>PRECAUTIONS:</u></p> <ul style="list-style-type: none"> • Monitor blood glucose. 	<p>MAX ADMINISTRATIONS: 3</p> <p><u>PRECAUTIONS:</u></p> <ul style="list-style-type: none"> • Admin. in 1h.